



Effect of Inhaled Tacrolimus on Ischemia Reperfusion Injury in Rat Lung Transplant Model

Bayer, Johanna¹ , Das, Nitin A.¹ , Baisden Clinton E.¹ , Rani, Meenakshi² , Schwacha, Martin G.², Sako Edward Y¹ , Williams, Robert O.,III⁴, Peters, Jay I.³, Johnson, Scott B.¹

¹Cardiothoracic Surgery, UTHSCSA, San Antonio, TX

²Trauma & Emergency Surgery, UTHSCSA, San Antonio, TX

³Division of Pulmonology, Department of Medicine, UTHSCSA, San Antonio, TX,

⁴Austin College of Pharmacy, Austin, TX




Disclosures

- Robert Williams and Jay Peters: Patent rights for inhaled tacrolimus have been sold to the UT Board of Regents.
- None of the other authors have anything to disclose



Problems

- Ischemia Reperfusion Injury (IR Injury) is inevitable in transplantation
- Contributes to primary graft dysfunction (PGD)
- Negatively affects overall survival after lung transplantation




Inhaled nanoparticle Tacrolimus

Standard




Diameter: 10 000 nm

Nanoparticle




Diameter: 200 nm

Pharmacokinetics



Watts AB, Saad AR, Johnson SB et al, Characterization and pharmacokinetic analysis of tacrolimus dispersion for nebulization in a lung transplant rodent model, International Journal of Pharmaceutics, 2010

Efficacy





Simmons JD, Johnson SB et al, The Efficacy of Inhaled Nanoparticle Tacrolimus in Preventing Rejection in an Orthotopic Rat Lung Transplant Model, Aust society presentation June 2011

Hypothesis

Administration of inhaled nanoparticle
Tacrolimus to the **donor** prior to procurement
will attenuate ischemia reperfusion injury after
lung transplantation

Orthotopic Rat Lung Transplant



Study design

Control Group:
(n=6)


Treatment Group:
(n=6)

Inhaled
Lactose



Donor
pretreatment

Inhaled
Lactose + NP
Tacrolimus


- **Functional assessment**
 - Arterial Blood Gas
- **Cellular changes**
 - Flow cytometry
- **Cytokines**
 - Luminex Assay
- **Tacrolimus levels**
 - Liquid chromatography/mass spectrometry




Results: Oxygenation



Results: Neutrophils




Left Lung CD 11b High + Cells vs Tacrolimus level




Results: Cytokines


TNF



IL-1b



IL-6



Conclusion

Inhaled nanoparticle Tacrolimus treatment of lung donors is associated with attenuation of ischemia-reperfusion injury on a functional level in lung transplantation

Thank you

- Funding: Grant from Institute for Integration of Medicine and Science (IIMS), Clinical and Translational Science Award (CTSA) grant - UL1RR025767.
- Acknowledgement:
 - the UTHSCSA Department of Epidemiology and Biostatistics and in particular Benjamin R. Ehler for his statistical analysis
 - the Core for Advanced Translational Technologies at UTHSCSA and Dr. Carolina Livi for Luminex service to perform cytokine measurements.
 - the Biological Psychiatry Analytical Laboratory and Dr. Martin Javors for their help with the Tacrolimus levels.